

THE CASCADE CAVER

PROPERTY OF
THE WINDY CITY GROTTO
LIBRARY

VOLUME II NUMBER 7
JULY 1963
OFFICIAL PUBLICATION OF THE
CASCADE GROTTO N. S. S.
SEATTLE WASHINGTON

COMING EVENTS

Monday July 8 1963

Regular Meeting
Don Hollidays
15611 157th Avenue South East
Lake Hills (see map)

NEW MOUNT ST HELENS CAVE GROUP VISITED by Bill Halliday

On Saturday, June 1, Jan Utterstrom and I scouted a cave area well to the north of the "usual" Mt St Helens caves, previously visited by Bill Reese and Jan. Accompanying us was Joan Crysdale, who achieved the dream of every novice caver--found a new cave.

The caves are in the rough, brushy area west and somewhat north of the St Helens Trail, and well to the north of the McBride Lake road. It would be extremely difficult to locate them without the assistance of someone familiar with the area. Because of the lack of time, no mapping or measuring was accomplished above or below ground.

The caves occur in sinks in a somewhat sinuous "line" in a lava flow with some rather steep slopes. We did not enter the lowest known cave of this group, located at the south end of a very large sink. Jan reported that it consists of a single chamber about 50 feet long, containing breakdown so dangerous he thinks it should not be entered again. 5 bats were present at the time of that visit. For the record, we list it as Breakdown Cave.

Moss Cave, the next north (upslope), is tiny but beautiful. Ten feet upslope, it is blocked by a red lava seal; twenty feet downslope from the entrance, by a black lava seal. Moss and small ferns grow in profusion in the twilight zone. A grylloblattid was collected beneath a rock in the entrance sink. Spiderwebs are present, and Jan found a bat here on his previous trip.

The next cave upslope was named Railroad Track Cave, because of its principal feature: two sets of thin lateral crusts which merge farther down its 100 foot course. Finally, the fused crusts merge to form a tube in tube which blocks farther penetration. This is an exceptional feature. The upper end of the cave is blocked by a black lava seal. Ferns and moss are also present here.

The next two caves upslope appear to be parts of a small distributory complex, now separated by lava seals and breakdown. Their entrances are about 75 feet apart, which is less than the length of the passage extending from Arch Cave toward Surprise Cave. The latter is the lower and the more complex of the two. A narrow pit, opened by Jan previously, leads to the main passage which extends northward about 100 feet, apparently west of Arch Cave.

NEW MT ST HELENS CAVE GROUP (continued)

The downflow lead extends a lesser distance, past multiple points of piracy of a red lava flow, to a spacious chamber into which two confluent small tubes lead from the northeast. Neither is penetrable for more than a few yards; one is blocked by a flow overlaying pahoehoe. Leading south from the eastern tube is the crawlway in talus into the "Sleeping Woman" Room, which we did not attempt.

Nearby Arch Cave was so termed for the beautiful little arch formed by a very short upper level at the entrance. Because of an overhanging drop of 15 - 20 feet into the upper end of this cave, we visited only the lower end, which consists of a single tube some hundred feet long, containing a magnificent red lava tongue. This cave probably lies a little east of Surprise Cave. It terminates by a black tumulus.

Some 200 feet farther north is a prominent entrance to a short cave with two levels leading north about 50 feet, and-according to Jan- about the same distance south. On the east wall of the entrance sink is an unusually fine example of a lava spring, almost in full daylight. (Them what says there ain't no such animal, please go see.) This was named Lava Spring Cave.

Still farther up the flow, well up a considerable slope, is a round sink about 20 feet in diameter and depth. Jan reported a short passage leading downslope.

At this point, we headed north, then west, seeking some sinks visible on aerial photos, but apparently we did not go far enough. While returning toward the McBride Lake road, which we struck about $\frac{1}{2}$ mile farther west, Joan found a small hole which led into a stoopway chamber about 15 feet in diameter followed by a crawlway some 50 feet long, relatively free of flow marks - not unexpected in this shallow small tube.

Photographs were also taken in the upper section of Little Red River Cave, where marks have been very drastically changed by recent logging and blowdown.

REINSTATED MEMBERS

Mrs Dina Lockyer
Queen Charlotte City
B.C. #3069
Canada

Note: Name change from Dina Turco
change of address: Alaska
Highway, Canada

Stephen Knutson
6904 S W Canyon Crest Dr.
Portland 1, Oregon

PLACE NAMES ON THE NORTHERN HALF OF VANCOUVER ISLAND

Lime Creek tributary of the Zeballos River, north of Nootka Island

Marble River long river: two lakes, Pat and Alice; east of Quartzine.

Limestone Island in Otter Cove, just east of Johnstone Strait and Discovery Passag

MOUNT ADAMS TRIP REPORT by Maryalice York

Ice Cave Campground was the base for Dr Don Holliday, Jim Montgomery, Van York, Luurt Niewenhuis, Maryalice York, Tom Dryden, and Wilma Dryden, who went on the scheduled trip over the June 22 - 23rd weekend.

Arriving at the campgrounds around 3:30 A.M., the group immediately went into the cave, where less ice was noted than had been seen on an August trip of two years ago.

Later Saturday morning, on the way to Dynamited Cave, a short excursion was made to Spearhead Cave.

Jim Montgomery was unable to make the entrance to Dynamited, so he spent the day exploring the surrounding area and talking to people about caves. Jim was told that three persons in one party were killed in Dynamited Cave as the result of improper and insufficient equipment.

Rope ladders were used at the 12 foot drop and at the 40 foot drop. Belaying was done as needed. The men were unable, due to lack of equipment, to safely make their way around the top of the 50 foot drop, and came back to the Bubble Room where the distaff members were waiting.

Don Holliday, Tom Dryden, and Van York made a trip into the passage under the 12 foot drop.

On Sunday, New Cave was entered. About 3/5ths of the way in, part of a Chiroptera skeleton was found. About 100 yards past this, about 20 rodent skeletons were found. It is believed that these rodents had tunneled down from above. Samples of yellow and white slime were taken from New Cave at the 3/5ths way mark. Don Holliday added water samples from both New Cave and Dynamited Cave to his collection for analysis. Many soda straw like formations were noted, ranging in length from one to seven inches, in New Cave.

Dry Creek cave was visited with members using two different exits. Don Holliday and Jim Montgomery followed the system above ground.

Many pictures were taken on the trip with the slide series project in mind.

A Forest Service Road was used on the return trip to Seattle, cutting the mileage to 183 versa 300, but little time was saved.

Forest Service Maps of the Mount Adams area, showing all corrections for 1963, have been furnished to the Grotto and will be available to members.

COPIES OF FEATURE ARTICLE AVAILABLE

Additional copies of this months feature article "Concerning The Clayish-Slimy Formations Termed Vermicular" are available to interested persons and agencies, no charge. (Please send a stamped, self-addressed envelope however.) So far as this Grotto has been able to determine, this is the first technical paper on this phenomenon, published in the United States. Special thanks must be given to Mr. Rodolfo Musco for his efforts to make this report available.

ICE CAVE TRIP REPORT By Bob Ashworth


On May 11th, 1963, two friends and I drove to Arnold Ice Cave where we spent the night. The next day, May 12th, we went down into Arnold in which recently, stairs have been built, and noted that since January 5, 1963, the ice has built up on the floor about three inches. Next we want to look for the caves that Steve Knutson reported in the April, 1963 issue of the Cascade Caver on the north slope of Lava Top Butte. We found a small ice cave on the north side of the butte, (about 200 feet long). The cave did not fit the description given by Steve Knutson in the Cascade Caver. There were two or three inches of melt water on the top of the ice and we found old animal bones in the cave. This could possibly be the same cave that Steve found or it could possibly be a different one.

Printer's note: Herman, the friendly multilith, that has been printing the Caver for the past several months, has gone off on a summer vacation, so you are just going to have to put up with this messey and misspelled mimeograph sheet for a month. JAF

1964 CONVENTION: NEW BRAUNFELS, TEXAS

The National Board of Directors, at the meeting last February, selected New Braunfels, Texas as the site for the 1964 Convention. This carries out the policy of meeting in the east and the west in alternate years. New Braunfels is located on the southeastern edge of the Balcones Escarpment of the Edwards Plateau about 30 miles from San Antonio and 50 miles from Austin. The population is 15,000 but hotel capacity is quite adequate as well as excellent campground facilities. They say, "numerous local caves" and promise several field trips including Natural Bridge Caverns, the states largest cave, and one to Bustamante, Mexico. Mr James Estes will be convention chairman. He is Sec-Tres of the Texas Speleological Association, and Secretary of the Abilene Grotto.

MAP TO DON HOLLIDAYS HOME FOR REGULAR MEETING



PIETRO PARENZAN

CONCERNING THE CLAYISH-SLIMY FORMATIONS TERMED VERMICULAR

International symposium of Speleology
Varenna (Lake of Como), Italy 3-6 October 1960

From Memoria V della Rassegna Speleologica Italiana, translated by
Rodolfo Musco and Francis X. Connelly, edited by William R. Halliday

* * * *

During recent speleological meetings certain scholars have pointed out characteristic clayish formations noticed on walls and ceilings of some caves and have formulated some hypotheses about their origins.

The subject seems interesting, not only because the genetic factors still are not completely known--and indubitably they must be many in relation to the variety of the formations about which we are speaking in their macroscopic aspects--but also because they are discovered in various caves. They have been found particularly in Italy, in southern caves, and also in coastal caves whose origins are connected with the sea (Zinzulusa Cave, Puglie, etc.)

These formations, improperly named "clayish vermicular formations" with respect to Kyrle's classifications ought to belong, in my opinion, to a third group: mixed, because although they are mostly allochthonous deposits, they have, at least in part, character of autochthonous deposits. In fact, Kyrle defines deposits in this way:

AUTOCHTHONOUS DEPOSITS: 1) of physical origin as waters of internal condensation; ice from freezing of waters of condensation; rocky sediments caused by landslip or by internal splitting, etc.; 2) of chemical origin as crystalline concretions, concretions of soldering, moon milk concretions; 3) of biological origin as veils of the subterranean bacterial activity; ferrobacteria; sulphobacteria, etc.

ALLOCHTHONOUS DEPOSITS: They include water coming from the outside: water of subterranean streams and lakes; snow and ice; water-borne gravels, sands, slime and clays (red earth, bauxites); pyroclastic material of hydric and aeolian transportation; sands and dust of aeolian deflation; coarser gravels and sediments of glacial transportation; vegetal and animal organic remains (coprolites, phosphatic earths, guano, ossiferous breccias), etc. (Anelli, 1959).

In fact, these formations are internal, with material of both internal and external origin, and the genetic factors also belong to both groups: external and internal.

Anelli, in one of his recent works (1957-59), in the chapter about fill deposits, did not mention these formations, not even among the "clayish-slimy concretions" and among deposits of internal degradation. He did not mention in any way the often-used words "leopard hide" and "vermicular formations".

Martel, in his description of the Grand Aven de Canjuers in France, pointed out for the first time some worm-like formations that he called "clayish ribbons".

Jeannel and Recovitzá pointed out similar formations in the Rumanian cave "Pestarea mare de la Soronista", expressing the opinion that those ought to be attributed to dropping of water on levi-gated rock walls.

Also Gortani mentioned formations of the same kind in the caves of the Marches in eastern Italy, and in the Postumia Caves, one of the largest caves in northern Italy near the Yugoslavian borders.

Geze, in Renault's report, mentioned vermicular formations in Rochefort cave in Mayenne, France.

Marchetti calls the vermicular formations "leopard hide". Perna and Pozzi describe types completely differing from place to place in the Grotta del Fiume. They, however, could not identify a common characteristic aspect of the phenomenon in relation to their location. Those authors described these formations: "They are clayish ribbons, reddish, brown or yellowish, containing much water and can be removed by a touch of a finger. They are often inter-linked while at other times they appear as rounded and isolated maculations."

Perna and Pozzi (1959) say that some particular aspects "give the idea of worm-eating," and that others ... "give the idea of tiger hide more than leopard hide".

In the Cave of Castel Lueghi (Istrian Peninsula), the same authors have found, as in the Cave of the Holy Virgin of Frassassi, in the Grotta Bella and in the Grotta Sulfurea, some vermicular formations ... "that would merit more study." They have also found similar deposits in the "Pont Niv" or "Native Bridge", the upper branch of the Cunardo Cave in the Province of Varese, in the Ladin Dolomites. "They look mostly like half-ellipsoid spots; about 1-1.50 cm. long, and are mostly located on concretionary deposits on walls and niches. In some points those recoverings are so thin that they cover exactly the form of the greatly eroded, sub-standing wall. Some spots are covered with a very thin veil of calcite...are always of a great interest because they are the only ones known in Lombardy and are among the very few of this kind found in Italy."

Jeannel and Recovitzá have defined a type of formation as "clayish vermiculations" ("vermiculures argileuses").

From a genetic point of view, Jeannel and Recovitz are of the opinion that these formations, in their different aspects, have to be considered due "to dropping of water on vertical walls"; but Perna and Pozzi do not think that this explanation is sufficient for those formations that are "on non-vertical walls or even on ceilings of tunnels, where drops of water have a tendency to drip instead of flowing along the walls." The same authors believe that Renault's hypothesis is more appropriate, if not sufficient. They believe that it is a residual product of decalcification deposited near networks of tiny fractures, and they compare these formations with those figures that appear on certain cement-manufactured articles that "...are constituted of calcium carbonate coming from the carbonation of free lime-mortar contained in cement that is left as a deposit in holes of retirement."

Formations of this type have been attributed by Marchetti (1950) also to sulphur water, but Perna and Pozzi point out that in the caves studied by them there is a lack of sulphur water. They postulate "many other factors, as humidity of atmosphere (vermiculations are found on walls covered with little drops of condensed water), air currents, variations of temperature, gas dissolved in the atmosphere, etc."

At this point there are three questions that arise from the examination of these formations: a) their nature, meaning their constitution; b) their morphological structure, that gives the characteristic various macroscopic aspects; c) their genesis.

Regarding their nature, all the formations of this kind observed by me in different southern caves are constituted neither of products of decalcification nor of any other chemico-genetic process, but simply of material transported from the overburden, from material that fills niches, crevices, stratifications, etc., often mixed with surface material transported by draughts, bat guano and sand and bits of debris carried by waters.

An exceptional field of research about this subject that I have the pleasure to put to the scholars' disposition, is represented by the Experimental Subterranean Biological Station of Naples, founded by me many years ago in the caves and holes under Naples. The "nursery station" is a room covered with "opus reticulatum" of volcanic tufa that was part of the "cavea" of a Roman Theater built about 20 centuries ago. Its walls and ceiling are almost entirely covered by clayish-slimy formations, of the kind known as "leopard hide". It is not a calcarean cave, and we cannot talk, then, of calcification.

The material of clayish-slimy formations, of a dark brown color, examined through the microscope, is of nothing else but heterogeneous material, containing the same elements contained in the ceiling earth. Also present are fungus elements, some protozoic cysts, etc. I also found, in a preparation, some hooks attributable to cestodes of a microtenia (*Echinococcus*).

For my observations which agree also with observations of similar material found in various natural caves, let us go back to the hypothesis of Jeannel and Recovitz. I found that on non-vertical surfaces and on the ceiling of caves, water can move without having necessarily to drop from the ceiling, depending on humidity, saturation of the overburden, condensation and surface variations.

Also, the heterogeneousness of materials that may contain elements of varying weight and thickness may be added to the different factors that determine the formation of clayish earthy of these clayish-slimy structures.

I would like to point out that I was able to reproduce some vague similar formations experimentally. I used a glass ceiling, covered with a mixture of earth, sand, cement and sawdust, and I produced specified vibrations as a substitute for the hypothetical natural effects of air. Also I allowed little veins of water to flow on the prepared glass and they made similar formations, even if--as is natural--they were not so clear-cut as those formed naturally, through long periods of time under the action of complex factors not entirely known. Nevertheless, among the factors that enter without doubt in the genesis of this phenomenon, "water droppings" have a primary importance, following Jeannel and Recovitz, because without them the other factors could not have a way to be developed.

The theories and hypothesis formulated about the genesis of these formations are four: biological (which cannot be true); as a deposit of transportation (including also aeolic deposits); as an in situ product of corrosion; and, finally, as a fossil deposit. In this last case this filling up ought to be the remains of an older and more complete filling of the cave. The described case of the Subterranean Station of Naples, however, breaks down this hypothesis, because from the time of its formation to the present time the local has never been filled up with clay or earth, nor invaded afterwards by water.

According to Renault, a formation "in situ" produced by corrosion of waters of infiltration which had perspired through microholes could explain some cases. But this could not be applied to vermiculations which are found on stalagmitic casts.

The nature of the substratum of these formations of pure or stalagmitic limestone should force us, following Renault, to investigate the existence of phenomena of substitution if the studied forms, as ribbons or spots, were not in contrast with this interpretation.

Seeing superficially the nature of the material and of the phenomenon, and considering the different opinions about the genesis of these formations, I believe that what we know better are the different aspects of this phenomenon. I think it is better to put together these aspects in a scheme of classification as it is usual to do in similar cases. In fact, these deposits have different forms.

In southern caves I could find the types mentioned by the different authors, in formations that go from macular formations to thick and bubbled ones and more or less thin and sketched ones; from the hieroglyphic-like formations to those which are dendritic, and from the interlinked linear form to the irregular reticulum, etc.

Renault has tried to tabulate different types of vermiculations of two families, as follows:

FAMILIES	FORMS OF CORROSION	FORMS OF DEPOSIT
I. Fluoform Vermiculations	a) vermiculations of corrosion of saharian rocks.	b) vermiculations of calcite.
II. Structural vermiculations	a) Stilolytes. b) Alveolar vermiculations	c) manganese dendrites. d) clayish vermiculations

The character of genetic order common to these formations consists, according to Renault, of a slow and reduced water alimentation.

Both corrosional and depositional forms show well the aspects of vermiculations, considering, however, also the formations opposed to these forms of caves--i.e., the vermicular and alveolar furrows frequent in calcareous surface rocks.

I desire, to present a classification merely for forms of cavern deposits that cannot be all included under the d) letter of Renault's tabulation, because not all of them seem "vermicular". In fact, macular or blistered formations frequently do not have a vermicular aspect. For this reason I believe that all these formations may be classified as follows:

1st group:

CLAYISH-SLIMY

MACULATIONS

- 1) Point-like maculae
- 2) Placoid maculae
- 3) Blistered maculae
- 4) Irregular, elipsoid or elongated maculae.

2nd group:

CLAYISH-SLIMY

VERMICULATIONS

- 5) Large vermiculations (leopard hide)
- 6) Linear, elongated, simple or interlacing vermiculations (tiger hide)
- 7) Hieroglyphic-like vermiculations
- 8) Dendrite-like vermiculations

This table (see figure), I think, is useful to classify all the forms known today in an ordered series.

To conclude, up to this time there have been reported and more or less adequately described 8 fundamental forms of spelean clayish-slimy formations, whose genesis still is not entirely understood.

Nevertheless, we can recognize that, at least in the cases examined up to now, they are derived from materials of internal transfer, most likely from dropping, flowing or percolating water, through joints, microfissures, etc. These formations contain all the elements that may be found in the ground overlying the cave or in materials of old filling of niches, clefts, etc., mixed, sometimes, with elements coming from guano.

Also considering the examined formations, we have to exclude any chemico-genetic factor, giving more value to merely physical factors. I do not think, however, that we have to exclude a biological factor "of maintenance", at least for those cases--perhaps scarce--in which constituent materials of clayish-slimy formations are within reach of biological elements (fungus, bacterial and protozoan forms). The different kinds of formations, following Prof. Parenzan's classification, found in the Oregon Caves are those number 2, 4, 5, 7 and 8.





1



2




3




4




5



6



7



8



THE CASCADE CAVER
1117 - 36th AVENUE EAST
SEATTLE 2, WASHINGTON



Non-delivery due to failure of sender to supply complete or correct address. Do not reuse this envelope.



Wendy City Gratto, N.S.S.
430 South Euclid
Oak Park, Ill.

- Moved, left no address
- No such number
- Moved, not forwardable
- Addressee unknown

115

177798

177798

